arXiv:1504.04631v2 [math.DS] 31 Jul 2015

A regularity result for the fractional Fokker-Planck equation with Ornstein-Uhlenbeck drift

Xiaoxia Xie[a,*], Jinqiao Duan[a], Xiaofan Li[a], Guangying Lv[b]

[a] *Department of Mathematics, Illinois Institute of Technology, Chicago, IL 60616, United States*
[b] *Institute of Contemporary Mathematics, Henan University, Kaifeng, Henan 475001, China*

Abstract

Despite there are numerous theoretical studies of stochastic differential equations with a symmetric α-stable Lévy noise, very few regularity results exist in the case of $0 < \alpha \leq 1$. In this paper, we study the fractional Fokker-Planck equation with Ornstein-Uhlenbeck drift, and prove that there exists a unique solution, which is C^∞ in space for $t > 0$ when $\alpha \in (0, 2]$.

Keywords: Fractional Laplacian operator, the isotropic α-stable Lévy process, fractional Fokker-Planck equation, Ornstein-Uhlenbeck drift.

---

1. Introduction

In this paper, we are interested in the regularity of solutions of the generalized Fokker-Plank equation for $x \in \mathbb{R}^d$ and $t > 0$

$$\partial_t u = \mathcal{I}[u] - \nabla \cdot (f(x)u), \quad u(0, x) = u_0(x), \tag{1.1}$$

where $\mathcal{I}$ is an operator defined in (2.1), $f(x) : \mathbb{R}^d \to \mathbb{R}^d$ in the drift term is a deterministic driving force, and u_0 in $L^\infty(\mathbb{R}^d)$ is nonnegative. We assume without loss of generality that $\int_{\mathbb{R}^d} u_0(x)dx = 1$. When $\mathcal{I}$ is the classical Laplacian, Eq (1.1) is the classical Fokker-Plank equation. If $f(\cdot)$ fulfills the usual assumptions of regularity needed in the Laplacian perturbation theory, it is well-known that the solution is smooth due to the regularizing effect of the parabolic operator. If $f(\cdot)$ is less regular, there is a series of papers by Lions, *et al.* devoted to vector field $f(\cdot)$ having bounded divergence and some Sobolev type regularity, and it is shown that the H^1 regularity in x is sharp for $f(x)$ being in Sobolev spaces instead of being Lipschitz or continuously bounded (see [5, 6, 12, 13]).

The Laplacian is the infinitesimal generator of the Brownian motion, hence the classical Fokker-Planck equation corresponds to a stochastic system with Gaussian noises. However, the noise is often non-Gaussian, and in this paper, we focus on the symmetric α-stable Lévy motion, for which the infinitesimal generator is the fractional Laplacian, denoted by $\Delta^{\alpha/2}$ with $\alpha \in (0, 2)$. Eq (1.1) with $\mathcal{I} = \Delta^{\alpha/2}$ is called the fractional Fokker-Plank equation (see Section 2 for details).

There are many results (see [4], [7]-[11], [14], [18]-[22], [24]-[26],[28] and references therein) on gradient perturbation of the fractional Laplacian operator in the frame work of regular vector field, i.e. the well posedness and regularity of the solution to the equation $u_t = \Delta^{\alpha/2}u - f \cdot \nabla u$ with $u_0 \in L^\infty(\mathbb{R}^d)$ and f satisfying some regularity assumptions. Depending on the value of α, there are three cases. First, $\alpha \in (0, 1)$ is known as the supercritical case because $\Delta^{\alpha/2}u$ is of lower order than $f \cdot \nabla u$. In [24], it is shown that the solution becomes differentiable with Hölder continuous derivatives, if $f \in C^{1-\alpha+\beta}$ for any $\beta \in (0, \alpha)$. With a slightly weaker regularity assumption on f, it is shown in [25] that u is Hölder continuous if $f \in C^{1-\alpha}$ in space. Second, in the critical case of $\alpha = 1$, Silvestre in [25] shows that u

---

[*] Corresponding author

Email addresses: `xxie12@iit.edu` (Xiaoxia Xie), `duan@iit.edu` (Jinqiao Duan), `lix@iit.edu` (Xiaofan Li), `gylvmaths@henu.edu.cn` (Guangying Lv)

becomes Hölder continuous for positive time if $f \in L^\infty(\mathbb{R}^d)$, while the authors in [8] and [22] achieved the same result by different approaches when f belongs to the BMO class and $\text{div} f = 0$. Third, when $\alpha \in (1,2)$, the term $\Delta^{\alpha/2}u$ is dominant, which is called the subcritical case, and better regularities have been proved under more general conditions. Actually, in the one-dimensional case ($d = 1$), Droniou, *et al.* in [10] show that the unique solution exists and $u \in C^\infty((0,\infty), \mathbb{R})$ for $f \cdot \nabla u$ replaced by $\partial_x(f(u))$. Their result is generalized to multidimensional cases in [19] with the term $H(t,x,u,\nabla u)$ instead of $\partial_x(f(u))$. Note that $f(u)$ and $H(t,x,u,\nabla u)$ must satisfy certain conditions in order for $\Delta^{\alpha/2}$ to be the dominate term. The estimate of the fractional heat kernel perturbed by gradient operator is obtained in [4] if f belongs to a Kato class. See also [9, 7, 2] for two-sided fractional heat kernel estimates.

However, for the vector field f lying in some Sobolev spaces instead of being uniformly bounded in x, the drift term could be stronger than the diffusion term $\mathcal{I}[u]$ even in the case of $\mathcal{I} = \Delta$. Hence it would not be possible to prove any regularity for u in general and there are relatively fewer papers. For the fractional Fokker-Plank equation, the existence of weak L^p-solution was recently obtained in [26], if $f(\cdot) \in L^q_{loc}(\mathbb{R}^d)$, where q is the conjugate exponent of p. The solution is also unique with further assumptions on $f(\cdot)$. However, there is no regularity result so far.

In the rest of the paper, we focus on the fractional Fokker-Plank equation with the Ornstein-Uhlenbeck drift, i.e., Eq (1.1) with $\mathcal{I} = \Delta^{\alpha/2}$ and $f(x) = -x$,

$$\partial_t u = \Delta^{\alpha/2}u - \nabla \cdot (-xu), \quad u(0,x) = u_0(x). \tag{1.2}$$

In this case, the unique weak L^p-solution exists by applying the results in [26]. However, we show that the unique solution to Eq (1.2) is smooth in this work. More precisely, the solution of (1.2) can be represented as

$$u(t,x) := (p * u_0)(x) = \int_{\mathbb{R}^d} p(t, x-y)u_0(y)dy, \tag{1.3}$$

where $p(t,x,y) = p(t,y-x)$ is the fundamental solution of (1.2) (or p is called the heat kernel of the fractional Laplacian perturbed by the Ornstein and Uhlenbeck drift). One can easily check that

$$p(t,x,y) = e^{dt}\widehat{p}\left(\frac{e^{\alpha t}-1}{\alpha}, e^t x, y\right), \tag{1.4}$$

where $\widehat{p}(t,x,y) = \widehat{p}(t,y-x)$ is fundamental solution to the fractional heat equation, i.e., $\partial_t u = \Delta^{\alpha/2}u$ (or $\widehat{p}$ is called the fractional heat kernel). It is well-known that $\widehat{p}(t,\cdot)$ is C^∞ for all $t > 0$ when $\alpha \in (0,2)$ (see [1]). Due to (1.4), the solution of (1.2) has the same regularity as that the fractional heat equation, as shown in the following theorem.

Theorem 1.1. *Let $u_0 \in L^\infty(\mathbb{R}^d)$ and be continuous almost everywhere. The function u defined in* (1.3) *is the solution to* (1.2). *Furthermore, for $T > 0$ or $T = \infty$,*

$$u(t,\cdot) \in C^\infty(\mathbb{R}^d), \quad \forall t \in (0,T]. \tag{1.5}$$

We remark that from the stochastic view point, the α-stable Ornstein-Uhlenbeck process was first studied in the case of $\alpha = 1$ in [17], and the sharp estimates of the mean first exist time from the ball was given for $0 < \alpha < 2$ in [20], and it is shown in [21] that the Harnack inequality holds if $1 \le \alpha < 2$ or $\alpha < 1 = d$. From the pseudo differential equations point of view, see [15] for the exponential decay in time of the solution to the steady state of the associated stationary equation, and [16] for the applications of the Lévy logarithmic Sobolev inequality to the study of the regularity of the solutions of the fractional heat equation and the asymptotic behavior of the Lévy-Ornstein-Uhlenbeck process. For numerical results, see [14]. Finally, we refer to [17, 27, 29, 30] for applications of the fractional Ornstein-Uhlenbeck process in physical sciences.

We point out that our result in Theorem 1.1 is a little surprising, especially for the supercritical case ($\alpha \in (0,1)$), because it implies that the fractional Laplacian operator $\Delta^{\alpha/2}$ has the smoothing effect like the Laplacian in the Ornstein-Uhlenbeck drift case. We wish to expand the result to include more general drift terms in further studies. The paper is organized as follows: in Section 2, we introduce some notations and definitions. And the proof of Theorem 1.1 will be given in Section 3.

2. Preliminaries

Let us briefly recall basic definitions of the isotropic α-stable Lévy operator and the potential theory of the fractional Laplacian (see[1, 2, 3, 23]).

The Lévy-Khinchine formula implies that there exists a symmetric semi-definite $d \times d$ constant matrix $\sigma = (\sigma_{i,j})$, a constant vector $b = (b_i) \in \mathbb{R}^d$ and a nonnegative measure ν on $\mathbb{R}^d$ such that the Lévy operator writes:

$$\mathcal{I}[u](x) = \langle \sigma \nabla u; \nabla u \rangle + \langle b; \nabla u \rangle + \int_{\mathbb{R}^d} (u(x+z) - u(x) - \nabla u(x) \cdot z \mathbb{1}_{\{|z|<1\}}) \nu(dz), \tag{2.1}$$

where $\langle \cdot ; \cdot \rangle$ is the inner product. The so-called Lévy measure ν can be singular at the origin and satisfies $\nu(\{0\}) = 0$ and $\displaystyle\int \min(1, |z|^2) \nu(dz) < \infty$ and the matrix σ characterizes the diffusion (or Gaussian) part of the operator (it can be null if $\sigma = 0$), while b and ν characterizes the drift part and the pure jump part, respectively. The characteristic exponent of the Lévy process is denoted by $\psi(\xi) = \langle \sigma \xi ; \xi \rangle + \langle b; \xi \rangle + a(\xi)$, where $a(\xi) = \displaystyle\int_{\mathbb{R}^d} [1 - \cos(\xi \cdot z)] \nu(dz)$ is associated with the pure jump part of the underlying process. The symmetric α-stable Lévy process refers to the case $b = 0$, $\sigma = 0$, and $\nu(dz) = \mathcal{A}_{d,\alpha} \dfrac{dz}{|z|^{d+\alpha}}$, where $\mathcal{A}_{d,\alpha}$ is so chosen that $a(\xi) = |\xi|^\alpha$ for $\xi \in \mathbb{R}^d$ (see [4] and references therein).

For $\alpha \in (0, 2)$ and $\psi \in C_c^\infty(\mathbb{R}^d)$, the fractional Laplacian is

$$\Delta^{\frac{\alpha}{2}} \phi(x) := A_{d,\alpha}, \lim_{\epsilon \to 0} \int_{|z|>\epsilon} \frac{\phi(x+z) - \phi(x)}{|z|^{d+\alpha}} dz. \tag{2.2}$$

Note that if $\alpha = 2$ the Lévy operator has no jump part, and $\mathcal{I}[\cdot]$ is the Laplacian operator in this case.

We let $\widehat{p}(t, \cdot)$ be the smooth real-valued function on $\mathbb{R}^d$ with Fourier transform

$$\int_{\mathbb{R}^d} \widehat{p}(t, z) e^{iz \cdot \xi} dz = e^{-t|\xi|^\alpha}, \quad t > 0, \;\; \xi \in \mathbb{R}^d. \tag{2.3}$$

It follows from (2.3) that $\widehat{p}$ is the heat kernel of the fractional Laplacian and it is well-known that for $t > 0$, $x, y \in \mathbb{R}^d$, $x \neq y$, the sharp estimate of $\widehat{p}(t, x)$ is

$$\widehat{p}(t, x, y) \approx \min \left(\frac{t}{|x-y|^{d+\alpha}}, t^{-d/\alpha} \right) \tag{2.4}$$

(see [4, 7, 10, 19]), where the notation $\approx$ means that either ratio of the sides is bounded by $C \in (0, \infty)$, and C does not depend on the variables shown, here t and x. The estimate of the first order derivative of $\widehat{p}(t, x)$ is was derived in [4, Lemma 5], and we extend it to get the estimate of the m-th order derivative of $\widehat{p}(t, x)$ for any $m \geq 0$ as follows

$$\partial_x^m \widehat{p}(t, x) \approx \sum_{n=0}^{n=\lfloor \frac{m}{2} \rfloor} C_n |x-y|^{m-2n} \min \left\{ \frac{t}{|x-y|^{d+\alpha+2(m-n)}}, t^{-\frac{d+2(m-n)}{\alpha}} \right\}, \tag{2.5}$$

where $\lfloor \frac{m}{2} \rfloor$ means the largest integer that is less than $\frac{m}{2}$.

3. Proof of Main Result

Proof. To prove Theorem 1.1, we first prove the following result: for $u_0 \in L^\infty(\mathbb{R}^d)$ and $u_0(x)$ be continuous almost everywhere. The function $u(t, x)$ defined as

$$u(t, x) = \int_{\mathbb{R}^d} \widehat{p}(t, y-x) u_0(y) dy \tag{3.1}$$

is the solution to $u_t = \Delta^{\alpha/2}u$ with $u(0,x) = u_0(x)$, and for $T > 0$ or $T = \infty$, we have

$$u(t,\cdot) \in C^\infty(\mathbb{R}^d), \quad \forall t \in (0,T]. \tag{3.2}$$

Firstly, we show that $u(t,\cdot) \in C^\infty(\mathbb{R}^d)$ for all $t \in (0,T]$. Since $u_0 \in L^\infty(\mathbb{R}^d)$, and $\widehat{p}(t,\cdot) \in L^1(\mathbb{R}^d)$ for $t > 0$, $\widehat{p}(t,\cdot) * u_0$ is well-defined. By the Young's inequality for the convolution in (3.1), we have

$$\forall (t,x) \in (0,\infty) \times \mathbb{R}^d, \quad \|\widehat{p}(t,\cdot) * u_0(x)\|_{L^1(\mathbb{R}^d)} \le \|u_0\|_{L^\infty(\mathbb{R}^d)}. \tag{3.3}$$

By the smoothness of $\widehat{p}(t,x,y)$, to show (3.2), it is sufficient to show that

$$|\nabla_x^m \widehat{p}(t,x-y)| \in L^1(\mathbb{R}^d), \quad \forall m \ge 0, t_0 < t < T, \text{ and } x,y \in \mathbb{R}^d.$$

Indeed, by (2.5), we have

$$\begin{aligned}
\int_{\mathbb{R}^d} |\nabla_x^m \widehat{p}(t,x-y)| dy \le& \sum_{n=0}^{n=\lfloor \frac{m}{2} \rfloor} \int_{\mathbb{R}^d} |x-y|^{m-2n} \min\left\{ \frac{t}{|x-y|^{d+\alpha+2(m-n)}}, t^{-\frac{d+2(m-n)}{\alpha}} \right\} dy \\
=& \sum_{n=0}^{n=\lfloor \frac{m}{2} \rfloor} \int_{B(x,r)} |x-y|^{m-2n} t^{-\frac{d+2(m-n)}{\alpha}} dy + \sum_{n=0}^{n=\lfloor \frac{m}{2} \rfloor} \int_{B(x,r)^c} \frac{t}{|x-y|^{d+\alpha+m}} dy \\
\le& \sum_{n=0}^{n=\lfloor \frac{m}{2} \rfloor} t_0^{-\frac{d+2(m-n)}{\alpha}} \int_{B(x,r)} |x-y|^{m-2n} dy + \sum_{n=0}^{n=\lfloor \frac{m}{2} \rfloor} T \int_{B(x,r)^c} \frac{1}{|x-y|^{d+\alpha+m}} dy \\
<& \infty.
\end{aligned}$$

Sine $u_0 \in L^\infty(\mathbb{R}^d)$ and $\widehat{p}(t,\cdot) \in C^\infty(\mathbb{R}^d)$, the theorem of continuity under the integral sign gives (3.2). The uniqueness of the solution $u(t,x)$ defined in (3.1) follows from the Fourier Transform. And it is clear that for x^0 at which $u_0(x)$ is continuous, we have

$$\lim_{t \to 0, x \to x_0} u(t,x) = u_0(x^0). \tag{3.4}$$

Indeed, fix $x^0 \in \mathbb{R}^d, \epsilon > 0$. Suppose $u_0(x)$ is continuous at x_0. Choose $\delta > 0$ such that

$$|u_0(y) - u(x^0)| < \epsilon \quad \text{if } |y - x^0| < \delta, \text{ and } y \in \mathbb{R}^d.$$

Then if $|x - x^0| < \frac{\delta}{2}$, we have

$$\begin{aligned}
|u(t,x) - u_0(x^0)| \le& \int_{\mathbb{R}^d} \widehat{p}(t,x-y)|u_0(y) - u_0(x^0)| dy \\
=& \int_{B(x^0,\delta)} \widehat{p}(t,x-y)|u_0(y) - u_0(x^0)| dy + \int_{B(x^0,\delta)^c} \widehat{p}(t,x-y)|u_0(y) - u_0(x^0)| dy \\
\le& \epsilon + 2\|u_0\|_{L^\infty} \int_{B(x^0,\delta)^c} \frac{t}{|x-y|^{d+\alpha}} dy.
\end{aligned}$$

In the second term, we have $|y - x^0| > \delta$. Hence

$$|y - x^0| \le |y - x| + |x^0 - x| < |y - x| + \frac{\delta}{2} \le |y - x| + \frac{1}{2}|y - x^0|.$$

Thus $|y - x| \ge \frac{1}{2}|y - x^0|$. Consequently, the second term in the last line tends to 0 as $t \to 0^+$. Hence, if $|x - x^0| < \frac{\delta}{2}$, and $t > 0$ is small enough, $|u(x,t) - u_0(x^0)| < 2\epsilon$.

Second, set $\tilde{t} = \frac{e^{\alpha t} - 1}{\alpha}$ and $\tilde{x} = e^t x$. By (1.4), we have

$$\nabla_x^m p(t,x,y) = e^{dt} e^{mt} \nabla_{\tilde{x}}^m \widehat{p}(\tilde{t}, \tilde{x}, y), \tag{3.5}$$

Hence by (2.5), we have

$$\begin{aligned}\int_{\mathbb{R}^d} |\nabla_x^m p(t,x-y)|dy &\le e^{(d+m)t} \sum_{n=0}^{n=\lfloor \frac{m}{2} \rfloor} \int_{\mathbb{R}^d} |\tilde{x}-y|^{m-2n} \min\left\{ \frac{\tilde{t}}{|\tilde{x}-y|^{d+\alpha+2(m-n)}}, \tilde{t}^{-\frac{d+2(m-n)}{\alpha}} \right\} dy \\ &= e^{(d+m)t} \left[\sum_{n=0}^{n=\lfloor \frac{m}{2} \rfloor} \int_{B(\tilde{x},r)} |\tilde{x}-y|^{m-2n} \tilde{t}^{-\frac{d+2(m-n)}{\alpha}} dy + \sum_{n=0}^{n=\lfloor \frac{m}{2} \rfloor} \int_{B(\tilde{x},r)^c} \frac{\tilde{t}}{|\tilde{x}-y|^{d+\alpha+m}} dy \right] \\ &\le e^{(d+m)t} \sum_{n=0}^{n=\lfloor \frac{m}{2} \rfloor} \tilde{t}^{-\frac{d+2(m-n)}{\alpha}} \int_{B(\tilde{x},r)} |\tilde{x}-y|^{m-2n} dy + \sum_{n=0}^{n=\lfloor \frac{m}{2} \rfloor} \tilde{t} \int_{B(\tilde{x},r)^c} \frac{1}{|\tilde{x}-y|^{d+\alpha+m}} dy \\ &< \infty \end{aligned}$$

for any $0 < t_0 < t < T$. Hence, the theorem follows. $\square$

Acknowledgement

This work was partly supported by ANL Grant 3J-30001-0025A.

References

References

[1] D.Applebaum, Lévy processes and stochastic calculus, Cambridge university press, 2009.

[2] R.M. Blumenthal, R.K. Getoor, Some theorems on stable processes, Trans. Amer. Math. Soc. 95 1960 263-273.

[3] J.Bliedtner, W.Hansen, Potential theory: an analytic and probabilistic approach to balayage, Springer Science & Business Media, 2012.

[4] K.Bogdan, T.Jakubowski, Estimates of heat kernel of fractional Laplacian perturbed by gradient operators, Comm. Math. Phys. 271 (2007), no. 1, 179-198.

[5] C. Le Bris, P.-L. Lions, Existence and uniqueness of solutions to Fokker-Planck type equations with irregular coefficients, Comm. Partial Differential Equations 33 (2008), no. 7-9, 1272-1317.

[6] C. Le Bris, P.-L. Lions, Renormalized solutions of some transport equations with partially $W^{1,1}$ velocities and applications, Annali di Matematica pura ed applicata 183, no. 1 (2004), 97-130.

[7] K. Bogdan, A. Stós, P. Sztonyk, Harnack inequality for stable processes on *d*-sets, Studia Math. 158 (2003), no. 2, 163-198.

[8] L. Caffarelli, L. Silvestre, Regularity theory for fully nonlinear integro-differential equations, Communications on Pure and Applied Mathematics 62, no. 5 (2009) 597-638.

[9] Z.-Q. Chen, T. Kumagai, Heat kernel estimates for stable-like processes on *d*-sets, Stoch. Process. Appl. 108 (2003), no. 1, 27-62.

[10] J. Droniou, T. Gallouët, J. Vovelle, Global solution and smoothing effect for a non-local regularization of a hyperbolic equation, J. Evol. Equ. 3 (2003), no. 3, 499-521.

[11] J. Droniou, C. Imbert, Fractal first-order partial differential equations, Arch. Ration. Mech. Anal. 182 (2006), no. 2, 299-331.

[12] R. J. DiPerna, P.-L. Lions, Ordinary differential equations, transport theory and Sobolev spaces, Invent. Math. 98 (1989), no. 3, 511-547.

[13] R.J. DiPerna, P.-L. Lions, On the Cauchy problem for Boltzmann equations: global existence and weak stability, Ann. of Math. (2) 130 (1989), no. 2, 321-366.

[14] T. Gao, J. Duan, X. Li, Fokker-Plank equations for stochastic dynamical systems with symmetric Lévy motions, submitted.

[15] I. Gentil, C. Imbert, The Lévy-Fokker-Planck equation: Φ-entropies and convergence to equilibrium, Asymptot. Anal. 59 (2008), no. 3-4, 125-138.

[16] I. Gentil, C. Imbert, Logarithmic Sobolev inequalities: regularizing effect of Lévy operators and asymptotic convergence in the Lévy-Fokker-Planck equation, Stochastics: An International Journal of Probability and Stochastics Processes 81, no. 3-4 (2009) 401-414.

[17] P. Garbaczewski, R. Olkiewicz, Ornstein-Uhlenbeck-Cauchy process, J. Math. Phys. 41 (2000), no. 10, 6843-6860.

[18] J. He, J. Duan, H. Gao, A nonlocal Fokker-Plank equation for non-Gaussian stochastic dynamical systems, To appear in *Applied Math. Lett.*, 2015.

[19] C. Imbert, A non-local regularization of first order Hamilton-Jacobi equations, J. Differential Equations 211 (2005), no. 1, 218-246.

[20] T. Jakubowski, The estimates of the mean first exit time from a ball for the α-stable Ornstein-Uhlenbeck processes. Stochastic Process, Appl. 117 (2007), no. 10, 1540-1560.

[21] T. Jakubowski, On Harnack inequality for α-stable Ornstein-Uhlenbeck processes, Math. Z. 258 (2008), no. 3, 609-628.

[22] A. Kiselev, F. L. Nazarov, Variation on a theme of Caffarelli and Vasseur, Journal of Mathematical Sciences 166, no. 1 (2010) 31-39.

[23] N. M. Landkof, Foundations of modern potential theory. Vol. 180. Springer, 1972.

[24] L. Silvestre, On the differentiability of the solution to an equation with drift and fractional diffusion, Indiana University Mathematical Journal. 61 (2012), no. 2, 557-584.

[25] L. Silvestre, Hölder estimates for advection fractional-diffusion equations, Annali della Scuola Normale Superiore di Pisa. Classe di Scienze. Accepted for publication.

[26] J. Wei, R. Tian, Well-posedness for the fractional Fokker-Plank equations, submitted.

[27] S. Jespersen, R. Metzler, H.C. Fogedby, Lévy flights in external force fields: Langevin and fractional Fokker-Planck equations and their solutions, Phys. Rev. E 59, 2736.

[28] A. Stós, Symmetric α-stable processes on d-sets, Bull. Polish Acad. Sci. Math. 48 (2000), no. 3, 237-245.

[29] W. A. Woyczyński, Lévy processes in the physical sciences, Lévy processes, 241-266, Birkhäuser Boston, Boston, MA, 2001.

[30] B.J. West, P. Grigolini, R. Metzler, T. F. Nonnenmacher, Fractional diffusion and Lévy stable processes, Phys. Rev. E (3) 55 (1997), no. 1, part A, 99106.